

CONTAGIOUS COMMENTS

Department of Epidemiology

Influenza 2006 – 2007: Vaccinate Children!

*Susan Dolan, MS, RN, CIC
Christine Robinson, PhD
Ann-Christine Nyquist, MD, MSPH*

What is influenza?

Influenza is a contagious respiratory disease caused by the influenza virus. Influenza is spread from person to person by direct contact, large droplet infection, or items recently contaminated by nasal pharyngeal secretions. The incubation period is short (usually 1 to 4 days) with an average of 2 days.

The illness in adults, older children and adolescents is characterized by an abrupt onset of fever, chills, myalgias, intense headache, and severe malaise accompanied by cough, sore throat and nasal congestion.

Illness can differ greatly in children. Neonates often present with a sepsis-like picture including lethargy, decreased eating and mottling. Infants and toddlers tend to present with gastrointestinal symptoms (nausea, vomiting and diarrhea), fever, anorexia and various respiratory syndromes. More severe illness can result if either primary influenza pneumonia or secondary bacterial pneumonia occurs.

Influenza virus subtypes are based on their surface hemagglutinin (H) or neuraminidase (N) proteins. Infection with one subtype offers little or no protection against virus of other subtypes. Repeated influenza epidemics persist because the type A and type B viruses undergo constant and rapid change due to antigenic drift. Major antigenic shifts occur with influenza type A and are the reasons for pandemics in susceptible populations. Critical to the propagation of an influenza pandemic is a new circulating influenza A strain, a susceptible population and most importantly human to human spread.

The trivalent influenza vaccine is updated annually to include viruses that have been in the worldwide circulation. The antigenic characteristics of current strains provide the basis for selecting which strain to include in each year's vaccine. When there is a good match between vaccine and circulating viruses, influenza vaccine has been shown to prevent illness in approximately 70–90% of healthy persons less than 65 years of age.

In the United States, two measures are available that can reduce the impact of influenza: immunoprophylaxis with vaccine (inactivated or live virus) and chemoprophylaxis or treatment with influenza specific antiviral drugs (i.e., amantadine, rimantadine, oseltamivir, and zanamivir).

What's New for the 2006-2007 Season:

1. ACIP has expanded its influenza vaccine recommendations to include children aged 6-59 months and their household contacts and out-of-home caregivers.

2. Ritzwoller, et. al., published data in *Pediatrics 2005* highlighting the importance of 2 doses of vaccine for previously unvaccinated children <9 years of age. Children who only received 1 dose of influenza vaccine showed no statistically significant reduction in influenza like illness (ILI) or pneumonia and influenza (P&I) when compared to unvaccinated children. Vaccine effectiveness in children who were fully vaccinated was 25% for ILI and 49 % for P&I.

3. Time interval between doses for those needing 2 doses varies depending on type of vaccine.

TIV (injectable)	4 weeks
LAIV(nasal)	6-10 weeks

4. Amantadine or rimantadine should not be used for treatment or chemoprophylaxis of influenza A in the U.S. due to widespread resistance of the influenza virus to these antiviral medications. Instead oseltamivir or zanamivir should be used for chemoprophylaxis treatment if indicated.

Vaccination Coverage for Kids Needs to Improve:

CDC recently reported that the influenza vaccine coverage rates for children 6-23 month in the US, nearly doubled between the 2003-04 season and the 2004-05 seasons. It is believed that the change from "encouragement" in 2003-04 to vaccinate this age group to a full recommendation in 2004-05 played an important role. Despite this increase, the percent of US children covered remains sub-optimal and only 17.8% were fully vaccinated. In Colorado, 40.4% received 1 or more doses of vaccine and 23.8% were fully vaccinated.

	1 or More Flu Dose	Fully Vaccinated
U.S.	33.4%	17.8%
Colorado	40.4%	23.8%
*Denver Area		25.2%

*Adams, Arapahoe, Denver and Douglas Counties

We know that children <2 years of age are at a high risk for hospitalization due to influenza and that children 24-59 months are more likely to seek medical care (ED, hospital, clinic). So,

the message is clear, vaccinate children, and their close contacts (household and daycare providers).

2006-2007 Influenza Season Vaccine:

Annual influenza vaccine is required to refresh immunity to strains of flu that may return. Vaccination each year is also recommended to allow for refinement of the vaccine to closer match strains that are in circulation in the current season.

This year influenza vaccine will contain antigen for the following strains:

- A/New Caledonia/20/99 (H1N1)-like
- A/Wisconsin/67/2005 (H3N2)-like (or the equivalent A/Hiroshima/52/2005)
- B/Malaysia/2506/2004-like (or the equivalent B/Ohio/1/2005 virus)

We anticipate a good supply (110 to 115 million doses) of influenza vaccine this year with the addition of 2 new vaccine manufacturers.

Influenza Vaccine Dosage by Age Group for the United States, 2006 – 2007 Season

Age Group	Dose	Number of Doses	Route	Product
6 – 35 months	0.25 mL	1 or *2	Intramuscular	Fluzone®
3 – 8 years	0.50 mL	1 or *2	Intramuscular	Fluzone® (or Fluvirin™ for 4 yrs of age and older)
More than 9 years	0.50 mL	1	Intramuscular	Fluzone® or Fluvirin™ (FLUARIX™ is only for 18 yrs and older)
Healthy people 5-8 yrs	0.5 mL	1 or **2	Intranasal (0.25 mL in each nostril)	FluMist™ (LAIV)
Healthy people 9-49 yrs	0.5 mL	1	Intranasal (0.25 mL in each nostril)	FluMist™ (LAIV)

* 2 doses a least 1 month apart if receiving vaccine for the first time

**2 doses at least 6 weeks apart if receiving vaccine for the first time

All U.S. inactivated influenza vaccine supply consists only of split virus vaccine. Because influenza vaccines have not been approved by FDA for use among children aged 0 – 5 months and because this group of children is at high risk for flu-related hospitalizations, vaccination is recommended for their household contacts and out-of-home caregivers.

What about the preservative vs. preservative-free vaccine?

A few years ago, the AAP made recommendations to decrease the exposure to thimerosal in young children (particularly less than 6 months of age) because of potential concern regarding excess mercury exposure. The only injectable vaccine that is preservative-free is Fluzone® by Sanofi Pasteur and it is intended for use in young children 6 to 35 months of age. TCH will have a supply of preservative-free vaccine this year. In addition, FluMist™ (LAIV) does not contain any preservative.

What about intranasal flu vaccine?

FluMist™ is approved for use in healthy (no underlying chronic medical condition) persons aged 5–49 years of age. It should not be given to pregnant women. FluMist™ is an intranasally

administered, trivalent, cold-adapted, live attenuated influenza vaccine (LAIV). The attenuated (weakened) virus is adapted to growing at colder temperatures, which means that the live virus vaccine grows in the cooler upper respiratory tract and stimulates immunity without causing disease in the warmer lungs. Children aged 5–8 years who have never received influenza vaccine need two doses of LAIV (6–10 weeks apart).

With the recipient in the upright position, approximately 0.25mL is sprayed into each nostril.

LAIV must be stored frozen (-15°C or colder), but may be thawed in a refrigerator and stored at 2-8°C (36-46°F) for no more than 60 hours prior to use.

TCH will have some FluMist™ available for eligible patients and staff. FluMist is not recommended for TCH staff that have close contact with severely immunosuppressed patients. Specifically, FluMist™ is not recommended for healthcare workers caring for patients in BMT, Hematology/Oncology, and solid organ transplant services. If a healthcare worker receives LAIV, the healthcare worker should refrain from contact with these severely immunosuppressed patients for 7 days after vaccine receipt.

Who should be vaccinated?

We recommend that the following people be vaccinated at the beginning of the 2006 – 2007 influenza season:

- All healthcare workers in hospitals and outpatient / community / homecare settings because they can transmit influenza to vulnerable patients, residents, etc.
- All children ages 6 – 59 months and their household contacts and out of home caregivers.
- All children and adults who have chronic disorders of the pulmonary or cardiovascular systems, including children with asthma.
- All children and adults who require regular medical visits due to chronic metabolic diseases, renal dysfunction, hemoglobinopathies, or immunosuppression, including persons with HIV.
- Children and teens (6 months – 18 years) who are receiving long-term aspirin therapy and might be at risk for Reyes syndrome after influenza infection.
- Household members (including siblings and out-of-home care providers) of persons in high-risk groups.
- Women who will be pregnant during the influenza season.
- Persons 50 years or older.
- Residents and employees of nursing homes and other chronic care facilities housing persons of age with chronic medical problems.
- Anyone wanting to decrease their risk of acquiring influenza infection this year.

What are the contraindications to vaccination?

- Hives or severe anaphylactic reaction to chicken, eggs or a previous flu vaccine.
- Acute febrile illness (wait till symptoms have abated).
- Infants less than 6 months of age.

- For inactivated influenza vaccine only: An individual with a hypersensitivity to thimerosal (common preservative found in contact lens solution). However, most patients do not develop reactions even when patch or intradermal tests for thimerosal indicate hypersensitivity. When reported, hypersensitivity to thimerosal has usually consisted of local, delayed-type hypersensitivity reactions.
- For intranasal influenza vaccine only:
 - An individual with long term health problems (heart or lung disease, asthma, kidney disease, metabolic disease (diabetes), anemia and other blood disorders.
 - Children or adolescents on long-term aspirin therapy
 - Pregnant women
 - Anyone with a h/o Guillain-Barré syndrome

Procedure to Provide Influenza Vaccine to Parents and Siblings of Patients (Household Contacts):

A limited amount of free influenza vaccine is available to household members of TCH inpatients and outpatients. Check Planet TCH under “quick links” for information on how to obtain vaccine or check if your unit/dept has a vaccine book for this.

TCH Staff Immunizations:

This year, TCH held a flu vaccine drill for targeted staff working with high risk patients. The process will be used to evaluate our ability to reach employees in the event of any future pandemic or disaster scenarios that would require mass vaccination and/or prophylaxis. After this 3 day campaign, free immunizations for staff (TCH, UPI, contract, students, etc.) will continue to be available in Employee Health Services. For questions, call EHS at 303-861-6577.

What are the side effects?

Inactivated injectable influenza vaccine contains only non-infectious (inactivated) viruses; it therefore cannot cause influenza. The most frequent side effect of vaccination reported by less than one third of vaccines is soreness at the vaccination site that lasts for up to 2 days. Fever, malaise, myalgia, and other systemic symptoms occur infrequently and most often affect persons who have had no prior exposure to the influenza virus antigens in the vaccine (e.g., young children). These reactions begin six to twelve hours after the vaccination and may persist for 1 to 2 days. The most common side effects associated with LAIV include nasal congestion, scratchy throat and cough. Symptomatic relief can be obtained by using non-aspirin containing analgesics. **Aspirin should not be used due to the association of Reyes Syndrome with influenza in children.** Immediate (presumably allergic) reactions (e.g., hives, angioedema, allergic asthma, and systemic anaphylaxis) occur rarely and probably result from hypersensitivity to some vaccine component – a majority of which are most likely related to residual egg protein.

Can influenza vaccine be administered with other childhood vaccines?

Children may receive inactivated influenza vaccine at the same time they receive other routine vaccinations including pertussis vaccine (DTP or DTaP). Influenza and pneumococcal vaccine may be administered at the same time using different sites without increasing side effects. However, influenza vaccine must be administered each year whereas pneumococcal vaccine is not. For LAIV, refer to package insert.

Can pregnant women be immunized?

Because of the increased risk for influenza-related complications, it is recommended that women who will be pregnant during the influenza season should be vaccinated with inactivated influenza vaccine. Vaccination can occur in any trimester. One study of influenza vaccination of more than 2,000 pregnant women demonstrated no adverse fetal effects associated with influenza vaccine.

TCH Employee Health will administer the vaccine to pregnant staff. **Reminder:** LAIV is contraindicated for pregnant women.

Note: Lactating Mothers.

“Inactivated influenza vaccine is safe for mothers who are breastfeeding and their infants. Because excretion of LAIV in human milk is unknown and because of the possibility of shedding vaccine virus given the close proximity of a nursing mother and her infant, caution should be exercised if LAIV is administered to nursing mothers. Breastfeeding does not adversely affect the immune response and is not a contraindication for vaccination”.

What is the appropriate isolation for influenza?

For children hospitalized with symptomatic or confirmed influenza, Droplet Precautions (mask, gown, and gloves) are recommended for the duration of the illness.

What about Avian Influenza?

The current seasonal influenza vaccine probably does not provide protection against the H5N1 avian influenza strain circulating in birds in Southeast Asia. At this time there is rare human to human transmission of this strain of influenza. No human illness caused by avian influenza has been seen in this country to date. Key components to prevention of influenza infection still hold: good hand hygiene, respiratory etiquette (cover your mouth and nose when you cough or sneeze), droplet precautions for patients in healthcare settings, influenza vaccination and laboratory based viral surveillance. Regularly updated information can be obtained on the CDC website <http://www.cdc.gov/flu/avian/> and <http://pandemicflu.gov/>.



Laboratory Testing

Christine Robinson, PhD

How does TCH test for influenza?

Two laboratory tests are available at TCH for influenza virus detection:

1. Respiratory virus DFA and Culture:

This test is performed by staining specimens with virus-specific monoclonal antibodies labeled with fluorescent compounds that cause infected cells to glow under a microscope if viral proteins are inside. Sensitivity is about 85% for influenza A and 75% for influenza B relative to culture. Two helpful features of the DFA are that the specimen adequacy can be determined, and 8 viruses (RSV, parainfluenza, influenza A, influenza B, adenovirus, and human metapneumovirus) can be detected simultaneously. The DFA is performed at least twice a day on weekdays, and once a day on weekends, year-round. It can be ordered one of three ways:

- As a stand-alone test for ED or short-stay hospitalized patients who will benefit only from rapid results;
- With backup culture if the DFA is negative for admitted patients who can benefit from more complete and sensitive results;
- With concurrent culture for very high risk patients (e.g., individuals with HIV or organ transplant) who need all viruses detected.

2. Influenza A+B IA:

A rapid, influenza immunoassay (IA) is available during “flu season.” This test is most useful to screen suspected “flu” cases seen in the ED late afternoons, evenings and nights when the respiratory virus DFA is not available. Unfortunately the IA will miss about 10% of pediatric DFA-positive influenza cases.

Order the flu IA when a rapid diagnosis of influenza A or influenza B will impact patient care, e.g., when decisions about administering antiviral medications or withholding antibiotics are being made. If the IA is negative and a more comprehensive and complete answer is needed, respiratory virus DFA can be performed on the same sample the following day. Results for the Flu IA are available in less than an hour any time of the day or night.



Chemoprophylaxis

1. Neuraminidase Inhibitor: (Influenza A or B Use):

Zanamivir (Relenza®) and oseltamivir (Tamiflu®) are approved for the chemoprophylaxis and treatment of influenza A and B. Treatment has been shown to decrease the duration of flu-related symptoms by 1 to 1.5 days. Oseltamivir has been approved for chemoprophylaxis and treatment of patients older than one year old. Zanamivir has been approved for treatment in patients 7 years and older and chemoprophylaxis of patients age 5 years and older.

Limitations of Both Medications:

- Must be administered within 48 hours of onset of symptoms.
- Not shown to prevent disease transmission.
- Have not been adequately studied in patients with serious health conditions, with renal or hepatic impairment.

a. Zanamivir (Relenza®):

Available as a dry powder administered via oral inhalation with a plastic device. The dose is two breath-activated inhalations (one 5mg blister per inhalation = 10mg) BID for 5 days.

Note: The product will be packaged in a foil pack (Roto Disk) containing 4 blisters of the drug. Five Roto Disks will be packaged in a tube (equals entire treatment course). The package also includes one Diskhaler device.

Contraindications / Precautions:

Zanamivir is not recommended for use in patients with underlying airway disease including asthma or COPD because of a lack of safety and efficacy data in these patients. **Serious adverse events including bronchospasm and decline in lung function have been reported with zanamivir use, most commonly in patients with underlying airway**

disease. (If zanamivir is used in patients with underlying airway disease, they should be instructed to have a fast-acting bronchodilator available.)

b. Oseltamivir (Tamiflu®):

Given twice daily for 5 days, with dose adjustments required in renal impairment. As with Zanamivir, Oseltamivir therapy should be initiated within 48 hours of onset of influenza symptoms.

Pediatric Dosing:

1 – 12 Years: 2mg/Kg/dose bid x 5 days (max. dose = 75mg).
13 Years & Older: 75mg bid x 5 days.

Additionally, oseltamivir has shown some benefit as a prophylactic agent for seasonal influenza when given once daily for 6 weeks, although the cost may be prohibitive. CDPHE strongly discourages “personal stockpiling” of Tamiflu (<http://www.cdph.state.co.us/dc/Influenza/avian/index.html>).

References:

CDC: Prevention and Control of Influenza: Recommendation of the Advisory Committee on Immunization Practices (ACIP). July 28, 2006 / 55(RR10) p1-42 <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5510a1.htm>

CDC. Childhood Influenza Vaccination Coverage – United States, 2004-05 Influenza Season. MMWR. October 6, 2006/Vol.55/No.39 p. 1061-5.

Bug Watch



Up-to-date information on currently circulating respiratory and enteric viruses and *B. pertussis* detected by the TCH Laboratory provided to you weekly. Posted on the TCH Internet <http://www.thechildrenshospital.org/pro/publications/bug.pdf> and/or sent by broadcast FAX. Contact Carolyn Brock by e-mail (brock.carolyn@tchden.org) or phone (303-861-6412) to begin receiving your personal copy.



The Children's Hospital Association
1056 E. 19th Ave., Denver, CO 80218-1088

CONTAGIOUS COMMENTS
Department of Epidemiology

NON-PROFIT
ORGANIZATION
U.S. POSTAGE
P A I D
DENVER, CO
PERMIT NO. 1050

EDITOR:

Barb Jensen, Staff Assistant III
The Children's Hospital, Dept. of Epidemiology, B-276
1056 E. 19th Avenue, Denver, CO 80218
Phone: 303-861-6072; FAX: 303-837-2631

jensen.barbara@tchden.org
<http://www.thechildrenshospital.org>

We Recycle!

©